

Ma 221

Chapter 1 - Basic Concepts

Classification of Differential Equations

A differential equation is an equation involving an unknown function and one or more of its derivatives. Thus it is a relation of the form

$$F\left(x, y, \frac{dy}{dx}, \dots, \frac{d^n y}{dx^n}\right) = 0$$

F is given, and we are to find y . The above is an ordinary differential equation of order n .

Example

$$\frac{dy}{dx} = f(x) \text{ or } y' = f(x)$$

Definition. The order of a differential equation is the order of the highest derivative appearing in the equation. If the equation is a polynomial in the unknown function and its derivatives, then the degree of such an equation is the power to which the highest derivative is raised.

Example $a(x)y'' + b(x)y' + c(x)y = f(x)$. second order, first degree

Remark: $f(x) = 0 \Rightarrow$ homogeneous equation. $f(x) \neq 0$ nonhomogeneous.

Example $\left(\frac{d^3 s}{dt^3}\right)^2 + 5s^4 t^3 = 0$ 3rd order, 2nd degree

Example $2xy'' - (x+3)y' + 6x^4y = 0$ 2nd order, first degree

Up until now we have mentioned only ordinary differential equations. We shall eventually be concerned with partial differential equations also.

Example

$$uu_{xx} = \frac{1}{c^2} u_t$$

Here $u = u(x, t)$. This is a partial differential equation. Here c is a constant.

Solutions of Differential Equations.

Consider the n -th order ordinary differential equation

$$F\left(x, y, \frac{dy}{dx}, \dots, \frac{d^n y}{dx^n}\right) = 0. \quad (1)$$

Definition. A solution of the ordinary differential equation (1) is a real valued function $y(x)$ defined on some interval I such that:

1. $y(x)$ and its first n derivatives exist for each $x \in I$.
2. The substitution of $y(x)$ into the differential equation makes the equation an identity in the interval I .

Remarks: I may be $(-\infty, \infty)$, $[a, b]$, (a, b) , $(a, b]$, $[a, b)$. We assume I is not degenerate. Now $1 \Rightarrow y$ and its $n-1$ first derivatives are continuous.

Example $y' = x$ $-\infty < x < \infty$ $y = \frac{1}{2} x^2$ is a solution since

$$\frac{dy}{dx} = \frac{d}{dx} \left(\frac{1}{2} x^2 \right) = x$$

Thus $y(x) = \frac{1}{2}x^2$ is solution for $-\infty < x < \infty$.

The function

$$y = \begin{cases} \frac{1}{2}x^2 - 1 & x \geq 0 \\ \frac{1}{2}x^2 & x < 0 \end{cases}$$

is not a solution of the differential equation on $-\infty < x < \infty$ due to the discontinuity at $x = 0$. $y = \frac{1}{2}x^2 - 1$ is a solution on $0 < x < \infty$ whereas $y = \frac{1}{2}x^2$ is a solution on $-\infty < x < 0$.

Example $y' + y = 0$ One solution is $y = e^{-x}$. The general solution is $y = ce^{-x}$, where c is any constant.

Remarks about solutions:

1. Sometimes we obtain the solution to a differential equation implicitly in the form $f(x, y) = 0$. We need not always solve for y as a function of x (cannot). However, we can verify that we have a solution by implicit differentiation.

Example $e^y \frac{dy}{dx} + x = 0$
 $\Rightarrow e^y dy + x dx = 0$
 $\Rightarrow e^y + \frac{x^2}{2} = c \quad (*)$

We could write $y = \ln(c - \frac{x^2}{2})$ but need not. To see if $(*)$ is a solution we differentiate implicitly. $(*)$
 $\Rightarrow e^y \frac{dy}{dx} + x = 0$.

2. Not all equations have solutions.

Example $(y')^2 + y^2 = -1$ has no solution.

Clearly $y = 0$ is not a solution. If $y \neq 0$, $\Rightarrow y^2 > 0$ and $(y')^2 > 0$.

Please present the material up to here.

Also, please present the solutions to problems 4 and 6 on page 14 of the text.

Initial and Boundary Value Problems

We have seen above that a differential equation need not have a unique solution.

Example $y' = x$ $y = \frac{1}{2}x^2 + c$.

If we are given some subsidiary condition then we will “pick” out a unique solution. For example, if we are given the initial conditions $y(0) = -1 \Rightarrow c = -1 \Rightarrow$ and $y = \frac{1}{2}x^2 - 1$.

For first order equations one is given one condition. For second order equations one needs two conditions.

Example $y'' + y = 0$

One may verify directly that $y = c_1 \sin x + c_2 \cos x$ is the solution, where c_1 and c_2 are constants.

If, for example, we are given $y(0) = 0$ and $y'(0) = 1 \Rightarrow$

$y(0) = c_1 \sin 0 + c_2 \cos 0 = c_2 = 0 \Rightarrow y = c_1 \sin x \Rightarrow y'(x) = c_1 \cos x \Rightarrow y'(0) = c_1 = 1$

Thus $y = \sin x$ is the solution.

We could have been given the boundary conditions $y(0) = 0$ $y(\frac{\pi}{2}) = 2$

$\Rightarrow c_2 = 0$ as before. Also $y(\frac{\pi}{2}) = c_1 \sin \frac{\pi}{2} = 2 \Rightarrow c_1 = 2. \Rightarrow y = 2 \sin x$

The above are two different kinds of conditions. When the two conditions are given at the *same* point,

they are called Initial Conditions; when the two conditions are given at two *different* points, they are called Boundary Conditions.

The equation together with the two conditions is called either an Initial Value Problem (I.V.P.) or a Boundary Value Problem (B.V. P.).

Example

$$\text{DE } y'' = 2x$$

$$\text{B.C. } y(0) = 0 \quad y(2) = 1$$

This is a B.V.P.

$$y' = x^2 + c_1$$

so

$$y = \frac{x^3}{3} + c_1x + c_2$$

$$y(0) = 0 \Rightarrow c_2 = 0 \quad y(2) = 1 \Rightarrow \frac{8}{3} + 2c_1 = 1 \Rightarrow 2c_1 = 1 - \frac{8}{3} = -\frac{5}{3} \text{ and therefore } c_1 = -\frac{5}{6}$$

$$\Rightarrow$$

$$y(x) = \frac{x^3}{3} - \frac{5}{6}x$$

is the solution.

Example

$$\text{D.E. } y'' = 2x$$

$$\text{I.C. } y(1) = 0 \quad y'(1) = -1$$

This is an I.V.P.

$$y' = x^2 + c_1$$

$$\text{so } y(x) = \frac{x^3}{3} + c_1x + c_2$$

$$y(1) = 0 \quad \Rightarrow \quad \frac{1}{3} + c_1 + c_2 = 0$$

$$y'(1) = -1 \quad \Rightarrow \quad 1 + c_1 = -1$$

$$\Rightarrow c_1 = -2 \quad \text{and} \quad \frac{1}{3} - 2 + c_2 = 0 \quad \Rightarrow c_2 = \frac{5}{3}$$

Thus

$$y(x) = \frac{x^3}{3} - 2x + \frac{5}{3}$$

is the solution.