

Assignment 5

This homework is due *Tuesday* Oct 1.

There are total 41 points in this assignment. 33 points is considered 100%. If you go over 33 points, you will get over 100% for this homework and it will count towards your course grade (up to 115%).

Collaboration is welcome. If you do collaborate, make sure to write/type your own paper *and give credit to your collaborators in your pledge*. Your solutions should contain full proofs. Bare answers will not earn you much.

This assignment covers section 3.1 in Bartle–Sherbert.

- (1) In this exercise you have to deliver specific inequalities from the definition of the convergent sequence. In each case below, find a number $K \in \mathbb{N}$ such that the corresponding inequality holds for all $n > K$. Give a *specific natural number* as your answer, for example $K = 1000$, or $K = 2 \cdot 10^7$, or $K = 139$, etc. (Not necessarily the smallest possible.)

You can (but you are discouraged to) use a calculator if you want to. However, 1) this problem can be done without using a calculator, 2) even if you do use one, your answers still should easily verifiable without one.

(a) [2pt] $\left| \frac{890534890.6451}{n} \right| < 0.00019011$

(b) [2pt] $\left| \frac{100-n}{n} - (-1) \right| < 0.0054352$,

(c) [3pt] $\left| \frac{200^{10}n+10^{100}}{n^2-10^{200}} \right| < 0.1$,

(d) [3pt] $\left| \frac{\cos(863n)}{\log n} \right| < 0.032432$,

(e) [4pt] (See example 3.1.11(d)) $|\sqrt[n]{n} - 1| < 0.01$. (*This item is due Oct 8.*)

- (2) REMINDER. Recall that a sequence $X = (x_n)$ in \mathbb{R} **does not** converge to $x \in \mathbb{R}$ if there is an $\varepsilon_0 > 0$ such that for any $K \in \mathbb{N}$ there is $n_0 > K$ such that following inequality holds: $|x - x_n| \geq \varepsilon_0$.

In each case below find a *real number* ε_0 that demonstrates that (x_n) does not converge to x .

(a) [2pt] $x_n = 1 + 0.1 \cdot (-1)^{n+1}$, $x = 1$,

(b) [2pt] $x_n = 1/n$, $x = 1/2013$,

— see next page —

- (3) REMINDER. Recall definition of a sequence in \mathbb{R} converging to an $x \in \mathbb{R}$:
 Let $X = (x_n)$ be a sequence in \mathbb{R} , let $x \in \mathbb{R}$. $X = (x_n)$ converges to x if $\forall \varepsilon > 0 \exists K \in \mathbb{N} \forall n > K, |x - x_n| < \varepsilon$.
- Below you can find (erroneous!) “definitions” of a sequence converging to x . In each case describe, exactly which sequences are “converging to x ” according to that “definition”.
- (a) [3pt] Let $X = (x_n)$ be a sequence in \mathbb{R} , let $x \in \mathbb{R}$. $X = (x_n)$ “converges to x ” if $\forall \varepsilon > 0 \forall K \in \mathbb{N} \forall n > K, |x - x_n| < \varepsilon$.
(If you are confused at this point, think of the problem this way: suppose for some sequence (x_n) and a number $x \in \mathbb{R}$ you know that statement (a) is true. What can you say about (x_n) ?)
- (b) [3pt] Let $X = (x_n)$ be a sequence in \mathbb{R} , let $x \in \mathbb{R}$. $X = (x_n)$ “converges to x ” if $\exists K \in \mathbb{N} \forall \varepsilon > 0 \forall n > K, |x - x_n| < \varepsilon$.
- (c) [3pt] Let $X = (x_n)$ be a sequence in \mathbb{R} , let $x \in \mathbb{R}$. $X = (x_n)$ converges to x if $\exists \varepsilon > 0 \exists K \in \mathbb{N} \forall n > K, |x - x_n| < \varepsilon$.
- (d) [4pt] Let $X = (x_n)$ be a sequence in \mathbb{R} , let $x \in \mathbb{R}$. $X = (x_n)$ “converges to x ” if $\forall \varepsilon > 0 \exists K \in \mathbb{N} \exists n > K, |x - x_n| < \varepsilon$.
- (4) (Exercise 3.1.8) Let (x_n) be a sequence in \mathbb{R} , let $x \in \mathbb{R}$.
- (a) [4pt] Prove that $\lim(x_n) = 0$ if and only if $\lim(|x_n|) = 0$.
- (b) [3pt] Prove that if (x_n) converges to x then $(|x_n|)$ converges to $|x|$.
- (c) [3pt] Give an example to show that the convergence of $(|x_n|)$ does not imply the convergence of (x_n) .