

## Assignment 8

This homework is due *Thursday* Oct 30.

There are total 23 points in this assignment. 20 points is considered 100%. If you go over 20 points, you will get over 100% for this homework and it will count towards your course grade (up to 115%).

Collaboration is welcome. If you do collaborate, make sure to write/type your own paper *and give credit to your collaborators in your pledge*. Your solutions should contain full proofs. Bare answers will not earn you much.

This assignment covers section 4.1–4.2 in Bartle–Sherbert.

- (1) (4.1.15) Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined by setting  $f(x) = x$  if  $x$  is rational, and  $f(x) = 0$  if  $x$  is irrational.
  - (a) [2pt] Show that  $f$  has limit at  $x = 0$  (*Hint*: you can use squeeze theorem).
  - (b) [2pt] Prove that if  $c \neq 0$ , then  $f$  does not have limit at  $c$ . (*Hint*: you can use sequential criterion.)
  
- (2) [3pt] (Theorem 4.2.4 for difference) *Using  $\varepsilon$ - $\delta$  definition*, prove that limit of functions preserves difference. That is, prove the following:  
 Let  $A \subseteq \mathbb{R}$ ,  $c \in \mathbb{R}$  be a cluster point of  $A$ , and  $f, g$  be functions on  $A$  to  $\mathbb{R}$ .  
 If  $\lim_{x \rightarrow c} f = L$ , and  $\lim_{x \rightarrow c} g = M$ , then  $\lim_{x \rightarrow c} f - g = L - M$ .
  
- (3) [4pt] Using arithmetic properties of limit, find the following limits.
  - (a)  $\lim_{x \rightarrow 1} \frac{x^{100} + 2}{x^{100} - 2}$ .
  - (b)  $\lim_{x \rightarrow 0} \frac{(x+1)^{20} - 1}{x}$ .
  - (c)  $\lim_{x \rightarrow c} \frac{(x-c+1)^2 - 1}{x-c}$ .
  
- (4) (a) [3pt] (4.2.5) Let  $f, g$  be defined on  $A \subseteq \mathbb{R}$  to  $\mathbb{R}$ , and let  $c$  be a cluster point of  $A$ . Suppose that  $f$  is bounded on a neighborhood of  $c$  and that  $\lim_{x \rightarrow c} g = 0$ . Prove that  $\lim_{x \rightarrow c} fg = 0$ .  
 Explain why Theorem 4.2.4 (Arithmetic Properties of Limit) cannot be used.
  - (b) [1pt] ( $\sim$ 4.2.11b) Determine whether  $\lim_{x \rightarrow 0} x \cos(1/x^2)$  exists in  $\mathbb{R}$ .
  
- (5) [4pt] (4.2.13) Functions  $f$  and  $g$  are defined on  $\mathbb{R}$  by  $f(x) = x + 1$  and  $g(x) = 2$  if  $x \neq 1$  and  $g(1) = 0$ .
  - (a) Find  $\lim_{x \rightarrow 1} g(f(x))$  and compare with the value of  $g(\lim_{x \rightarrow 1} f(x))$ .
  - (b) Find  $\lim_{x \rightarrow 1} f(g(x))$  and compare with the value of  $f(\lim_{x \rightarrow 1} g(x))$ .
  
- (6) [4pt] (4.2.15) Let  $A \subseteq \mathbb{R}$ , let  $f : A \rightarrow \mathbb{R}$ , and let  $c \in \mathbb{R}$  be a cluster point of  $A$ . In addition, suppose  $f(x) \geq 0$  for all  $x \in A$ , and let  $\sqrt{f}$  be the function defined for  $x \in A$  by  $(\sqrt{f})(x) = \sqrt{f(x)}$ . If  $\lim_{x \rightarrow c} f$  exists, prove that  $\lim_{x \rightarrow c} \sqrt{f} = \sqrt{\lim_{x \rightarrow c} f}$ . (*Hint*:  $a^2 - b^2 = (a - b)(a + b)$ . Another hint: you will probably have to consider cases  $\lim_{x \rightarrow c} f = 0$  and  $\lim_{x \rightarrow c} f \neq 0$  separately.)