

**Assignment 1.**

This homework is due *Thursday*, September 10.

Collaboration is welcome. If you do collaborate, make sure to write/type your own paper and *credit your collaborators*. Your solutions should contain full proofs. Bare answers will not earn you much.

**1. QUICK CHEAT-SHEET**

REMINDER. On the set  $\mathbb{R}$  of real numbers there two binary operations, denoted by  $+$  and  $\cdot$  and called addition and multiplication, respectively. These operations satisfy the following properties:

- (A1)  $(a + b) + c = a + (b + c)$  for all  $a, b, c \in \mathbb{R}$ ,
- (A2)  $a + b = b + a$  for all  $a, b \in \mathbb{R}$ ,
- (A3) there exists  $0 \in \mathbb{R}$  s.t.  $0 + a = a + 0 = a$  for all  $a \in \mathbb{R}$ ,
- (A4) for each  $a \in \mathbb{R}$  there exists an element  $-a$  s.t.  $a + (-a) = (-a) + a = 0$ ,
- (M1)  $(ab)c = a(bc)$  for all  $a, b, c \in \mathbb{R}$ ,
- (M2)  $ab = ba$  for all  $a, b \in \mathbb{R}$ ,
- (M3) there exists  $1 \in \mathbb{R}$  s.t.  $1 \cdot a = a \cdot 1 = a$  for all  $a \in \mathbb{R}$ ,
- (M4) for each  $a \neq 0$  in  $\mathbb{R}$  there exists an element  $1/a$  s.t.  $a \cdot (1/a) = (1/a) \cdot a = 1$ ,
- (D)  $a(b + c) = ab + ac$  and  $(b + c)a = ba + ca$  for all  $a, b, c \in \mathbb{R}$ .
- (NT)  $1 \neq 0$ .

REMINDER. Let  $\mathbb{A}$  be a set with two operations  $+$  and  $\cdot$  satisfying A1–A4, M1–M3 and D, NT. (For example,  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}$ .) The set  $\mathcal{P} \subset \mathbb{A}$  is called the set of *positive elements* if

- (P1) If  $a, b \in \mathcal{P}$ , then  $a + b \in \mathcal{P}$  and  $ab \in \mathcal{P}$ ,
- (P2) If  $a \in \mathbb{A}$ , then exactly one of the following holds:  $a \in \mathcal{P}$ ,  $a = 0$ ,  $-a \in \mathcal{P}$ .

Then we say  $a < b$  if and only if  $b - a \in \mathcal{P}$ ;  $a \leq b$  if and only if  $b - a \in \mathcal{P} \cup 0$ .

**2. EXERCISES**

- (1) (Exercise 1.1.1 in Royden–Fitzpatrick) Let  $a, b \in \mathbb{R}$ . For  $a \neq 0$  and  $b \neq 0$ , show that  $(ab)^{-1} = a^{-1}b^{-1}$ . (Hint: check that  $a^{-1}b^{-1}$  satisfies definition of  $(ab)^{-1}$ .)
- (2) Show that  $-1 \cdot a = -a$  for all  $a \in \mathbb{R}$ .
- (3) (1.1.2) Verify the following:
  - (a) For each real number  $a \neq 0$ ,  $a^2 > 0$ . In particular,  $1 > 0$  since  $1 \neq 0$  and  $1 = 1^2$ .
  - (b) For each positive number  $a$ , its multiplicative inverse  $a^{-1}$  also is positive.
  - (c) If  $a > b$  for real numbers  $a, b$ , then
 
$$ac > bc \text{ if } c > 0 \text{ and } ac < bc \text{ if } c < 0.$$
 (Hint: determine whether  $ac - bc \in \mathcal{P}$ .)
  - (d) If  $a > b > 0$  and  $c \geq d > 0$  for real numbers  $a, b, c, d$ , then  $ac > bd$ .

— see next page —

- (4) In each case below, determine if  $P$  is a set of positive elements (i.e. if  $P$  satisfies P1–P2).
- (a)  $\mathbb{A} = \mathbb{Z}$ ,  $P = \mathbb{N}$ ,
  - (b)  $\mathbb{A} = \mathbb{Z}$ ,  $P = -\mathbb{N}$ ,
  - (c)  $\mathbb{A} = \mathbb{Q}$ ,  $P = \{r \in \mathbb{Q} : r > 1\}$ ,
  - (d)  $\mathbb{A} = \mathbb{C}$ ,  $P = \{z = x + iy \in \mathbb{C} : x > 0\}$ ,
  - (e) Prove that for  $\mathbb{A} = \mathbb{C}$ , there is no set of positive elements. (In other words, one cannot endow  $\mathbb{C}$  with a meaningful order.)
- (5) (1.1.4) Let  $a, b$  be real numbers.
- (a) Show that if  $ab = 0$  then  $a = 0$  or  $b = 0$ . (Hint: multiply  $ab$  by  $a^{-1}$ .)
  - (b) Verify that  $a^2 - b^2 = (a - b)(a + b)$  and conclude that from part (a) that if  $a^2 = b^2$ , then  $a = b$  or  $a = -b$ .
  - (c) Let  $c$  be a positive real number. Define  $E = \{x \in \mathbb{R} \mid x^2 < c\}$ . Verify that  $E$  is nonempty and bounded above. Define  $x_0 = \sup E$ . Show that  $x_0^2 = c$  by doing the following:
    - (i) Assume  $x_0^2 < c$ . Show that you can find  $\varepsilon > 0$  such that  $(x_0 + \varepsilon)^2$  is still  $< c$ . Explain what is wrong with that.
    - (ii) Assume  $x_0^2 > c$ . Show that you can find  $\varepsilon > 0$  such that  $(x_0 - \varepsilon)^2$  is still  $> c$ . Explain what is wrong with that.
 Use part (b) to show that there is a unique  $x > 0$  for which  $x^2 = c$ . It is denoted  $\sqrt{c}$ .
- (6) (1.2.12) Problem 5c (together with in-class proposition about  $\sqrt{2}$ ) prove existence of at least one irrational number (*irrational* means “real but not rational”). Granted that at least one irrational number exists, prove that irrational numbers are dense in  $\mathbb{R}$ .