

**Assignment 3.**

## Möbius transformations. Complex differentiation

This assignment is due Wednesday, Feb 11. Collaboration is welcome. If you do collaborate, make sure to write/type your own paper.

- (1) Find Möbius transformation that carries points  $-1, i, 1 + i$  into  
 (a)  $0, 2i, 1 - i$ , (*Hint*: From  $-1 \rightarrow 0$  you know that it has the form  $\frac{z+1}{az+b}$ .)  
 (b)  $i, \infty, 1$ . (*Hint*: As above, note that  $i \rightarrow \infty$ .)
- (2) Find the images of the following domains under the indicated Möbius transformations:  
 (a) The quadrant  $x > 0, y > 0$  if  $w = \frac{z-i}{z+i}$ .  
 (b) The half-disc  $|z| < 1, \text{Im } z > 0$  if  $w = \frac{2z-i}{2+iz}$ .  
 (c) The strip  $0 < x < 1$  if  $w = \frac{z}{z-1}$ .  
 (d) The strip  $0 < x < 1$  if  $w = \frac{z-1}{z-2}$ .  
 (*Hint*: You mainly need to keep track of the borders. It should help a bit to keep in mind that under  $z \rightarrow \frac{1}{z-b} + b$ , straight lines that do not pass through  $b$  go to circles that pass through  $b$ ; and straight lines that pass through  $b$ , go to straight lines that pass through  $b$ .)

- (3) Show that the function  $f(z) = z\text{Re } z$  is differentiable only at the point  $z = 0$ , and find  $f'(0)$ .
- (4) Find  $v(x, y)$  such that the function  $f(z) = 2xy + iv(x, y)$  is complex differentiable. Express  $f$  as a function of  $z$ .
- (5) Show that in polar coordinates, at every nonzero point of  $\mathbb{C}$ , Cauchy–Riemann equations take form

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \varphi}, \quad \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \varphi}.$$

(*Hint*: Differentiate  $\frac{\partial u(x, y)}{\partial r} = \frac{\partial u(r \cos \varphi, r \sin \varphi)}{\partial r}$  using chain rule. Do the same with other partial derivations  $\frac{\partial u}{\partial \varphi}, \frac{\partial v}{\partial r}, \frac{\partial v}{\partial \varphi}$ . Use “usual” Cauchy–Riemann equations.)

- (6) Let  $z_0 \neq 0$  and let  $f(z) = \ln r + i\varphi$ , where  $r = |z|$ ,  $\varphi \in \text{Arg } z$ , and  $\varphi$  is chosen so that  $f$  is continuous in a neighborhood of  $z_0$ . Prove that  $f$  is differentiable in a neighborhood of  $z_0$ .
- (7) Find an angle by which tangents to curves at  $z_0$  are rotated under the mapping  $w = z^2$  if  
 (a)  $z_0 = i$ , (b)  $z_0 = -1/4$ , (c)  $z_0 = 1 + i$ .  
 Also find the corresponding values of magnification.
- (8) Which part of the plane is shrunk and which part stretched under the following maps: (a)  $w = z^2$ , (b)  $w = z^2 + 2z$ , (c)  $w = 1/z$ ? (*Hint*: Whether a map  $f$  shrinks or stretches at  $z_0$  depends on  $|f'(z_0)|$ .)