

Assignment 3.

Möbius transformations. Complex differentiation

This assignment is due Wednesday, Feb 10. Collaboration is welcome. If you do collaborate, make sure to write/type your own paper.

- (1) Find Möbius transformation that carries points $-1, i, 1 + i$ into
 - (a) $0, 2i, 1 - i$, (*Hint*: From $-1 \rightarrow 0$ you know that it has the form $\frac{z+1}{az+b}$.)
 - (b) $i, \infty, 1$. (*Hint*: As above, note that $i \rightarrow \infty$.)
- (2) Find the images of the following domains under the indicated Möbius transformations:
 - (a) The quadrant $x > 0, y > 0$ if $w = \frac{z-i}{z+i}$.
 - (b) The half-disc $|z| < 1, \text{Im } z > 0$ if $w = \frac{2z-i}{2+iz}$.
 - (c) The strip $0 < x < 1$ if $w = \frac{z}{z-1}$.
 - (d) The strip $0 < x < 1$ if $w = \frac{z-1}{z-2}$.

(*Hint*: You mainly need to keep track of the borders. It should help a bit to keep in mind that under $z \rightarrow \frac{1}{z-b} + b$, straight lines that do not pass through b go to circles that pass through b ; and straight lines that pass through b , go to straight lines that pass through b .)

- (3) Show that the function $f(z) = z\text{Re } z$ is differentiable only at the point $z = 0$, and find $f'(0)$.
- (4) Find $v(x, y)$ such that the function $f(z) = 2xy + iv(x, y)$ is complex differentiable. Express f as a function of z .
- (5) Show that in polar coordinates, at every nonzero point of \mathbb{C} , Cauchy–Riemann equations take form

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \varphi}, \quad \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \varphi}.$$

(*Hint*: Differentiate $\frac{\partial u(x, y)}{\partial r} = \frac{\partial u(r \cos \varphi, r \sin \varphi)}{\partial r}$ using chain rule. Do the same with other partial derivations $\frac{\partial u}{\partial \varphi}, \frac{\partial v}{\partial r}, \frac{\partial v}{\partial \varphi}$. Use “usual” Cauchy–Riemann equations.)

- (6) Let $z_0 \neq 0$ and let $f(z) = \ln r + i\varphi$, where $r = |z|$, $\varphi \in \text{Arg } z$, and φ is chosen so that f is continuous in a neighborhood of z_0 . Prove that f is differentiable in a neighborhood of z_0 .
- (7) Find an angle by which tangents to curves at z_0 are rotated under the mapping $w = z^2$ if
 - (a) $z_0 = i$, (b) $z_0 = -1/4$, (c) $z_0 = 1 + i$.

Also find the corresponding values of magnification.
- (8) Which part of the plane is shrunk and which part stretched under the following maps: (a) $w = z^2$, (b) $w = z^2 + 2z$, (c) $w = 1/z$? (*Hint*: Whether a map f shrinks or stretches at z_0 depends on $|f'(z_0)|$.)