

Name: \_\_\_\_\_

Lecture Section \_\_\_\_

**Ma 221**

**Exam III B Solutions  
07S**

I pledge my honor that I have abided by the Stevens Honor System.

---

**You may not use a calculator, cell phone, or computer while taking this exam. All work must be shown to obtain full credit. Credit will not be given for work not reasonably supported. When you finish, be sure to sign the pledge.**

Score on Problem #1 \_\_\_\_\_

#2 \_\_\_\_\_

#3 \_\_\_\_\_

#4 \_\_\_\_\_

Total Score \_\_\_\_\_

**Note: A table of Laplace Transforms is given at the end of the exam.**

Name: \_\_\_\_\_

Lecture Section \_\_\_\_

**1 (25 pts.)** Use Laplace Transforms to solve

$$y'' + 4y = 0; \quad y(0) = 1, \quad y'(0) = 2$$

Solutions:

$$s^2 \mathcal{L}\{y\} - sy(0) - y'(0) + 4\mathcal{L}\{y\} = 0$$

Thus

$$s^2 \mathcal{L}\{y\} - s - 2 + 4\mathcal{L}\{y\} = 0$$

So

$$\mathcal{L}\{y\} = \frac{s+2}{s^2+4}$$

Therefore

$$\begin{aligned} y(t) &= \mathcal{L}^{-1}\left(\frac{s+2}{s^2+4}\right) = \mathcal{L}^{-1}\left(\frac{s}{s^2+4}\right) + \mathcal{L}^{-1}\left(\frac{2}{s^2+4}\right) \\ &= \cos 2t + \sin 2t \end{aligned}$$

**2a (10 pts.)**  $\mathcal{L}^{-1}\left\{\frac{2}{(s+1)(s^2+1)}\right\}.$

Solution:

Method 1, without complex variables:

$$\frac{2}{(s+1)(s^2+1)} = \frac{A}{s+1} + \frac{Bs+C}{s^2+1}$$

Multiplying by  $s+1$  and setting  $s = -1$ , yields

$$1 = A$$

Thus

$$\frac{2}{(s+1)(s^2+1)} = \frac{1}{s+1} + \frac{Bs+C}{s^2+1}$$

Setting  $s = 0$  yields

$$\frac{2}{1} = \frac{1}{1} + \frac{C}{1}$$

so

$$C = 1$$

and we have

$$\frac{2}{(s+1)(s^2+1)} = \frac{1}{s+1} + \frac{Bs+1}{s^2+1}$$

Setting  $s = 1$ , yields

$$\frac{2}{2(2)} = \frac{1}{2} + \frac{B+1}{2}$$

Thus

$$B+1 = 0$$

so

$$B = -1$$

Hence

Name: \_\_\_\_\_

Lecture Section \_\_\_\_

$$\frac{2}{(s+1)(s^2+1)} = \frac{1}{s+1} + \frac{-s+1}{s^2+1}$$

Therefore

$$\begin{aligned}\mathcal{L}^{-1}\left\{\frac{2}{(s+1)(s^2+1)}\right\} &= \mathcal{L}^{-1}\left\{\frac{1}{s+1}\right\} - \mathcal{L}^{-1}\left\{\frac{s}{s^2+1}\right\} + \mathcal{L}^{-1}\left\{\frac{1}{s^2+1}\right\} \\ &= e^{-t} - \cos t + \sin t\end{aligned}$$

Method 2, using complex variables:

$$\frac{2}{(s+1)(s^2+1)} = \frac{2}{(s+1)(s+i)(s-i)} = \frac{A}{s+1} + \frac{B}{s+i} + \frac{C}{s-i}$$

$s = -1$  gives  $A = 1$  as above.  $s = -i$  gives

$$\frac{2}{(-i+1)(-2i)} = \frac{2}{2(-1-i)} = \frac{-2}{2(1+i)} \times \frac{1-i}{1-i} = -\left(\frac{2-2i}{4}\right) = -\frac{1}{2} + \frac{1}{2}i = B$$

and  $s = i$  gives

$$\frac{2}{(i+1)(2i)} = -\frac{2}{2(1-i)} \times \frac{1+i}{1+i} = -\frac{1+i}{2} = -\frac{1}{2} - \frac{1}{2}i = C$$

Thus

$$\frac{2}{(s+1)(s^2+1)} = \left(\frac{1}{s+1}\right) + \left(-\frac{1}{2} + \frac{1}{2}i\right)\left(\frac{1}{s+i}\right) + \left(-\frac{1}{2} - \frac{1}{2}i\right)\left(\frac{1}{s-i}\right)$$

So

$$\begin{aligned}\mathcal{L}^{-1}\left\{\frac{2}{(s+1)(s^2+1)}\right\} &= \mathcal{L}^{-1}\left\{\frac{1}{s+1}\right\} + \left(-\frac{1}{2} + \frac{1}{2}i\right)\mathcal{L}^{-1}\left\{\frac{1}{s+i}\right\} + \left(-\frac{1}{2} - \frac{1}{2}i\right)\mathcal{L}^{-1}\left\{\frac{1}{s-i}\right\} \\ &= e^{-t} + \left(-\frac{1}{2} + \frac{1}{2}i\right)e^{-it} + \left(-\frac{1}{2} - \frac{1}{2}i\right)e^{it} \\ &= e^{-t} + \left(-\frac{1}{2} + \frac{1}{2}i\right)(\cos t - i \sin t) + \left(-\frac{1}{2} - \frac{1}{2}i\right)(\cos t + i \sin t) \\ &= e^{-t} - \cos t + \sin t\end{aligned}$$

SNB check  $\frac{2}{(s+1)(s^2+1)}$ , It's the Laplace transform of  $\sin t - \cos t + e^{-t}$

**2b (15 pts.)** Find  $\mathcal{L}^{-1}\left\{\frac{s}{s^2-3s+3}\right\}$ .

Solution

$$\begin{aligned}
\mathcal{L}^{-1}\left\{\frac{s}{s^2 - 3s + 3}\right\} &= \mathcal{L}^{-1}\left\{\frac{s}{s^2 - 3s + \frac{9}{4} + 3 - \frac{9}{4}}\right\} \\
&= \mathcal{L}^{-1}\left\{\frac{s}{(s - \frac{3}{2})^2 + \frac{3}{4}}\right\} \\
&= \mathcal{L}^{-1}\left\{\frac{s}{(s - \frac{3}{2})^2 + \left(\frac{\sqrt{3}}{2}\right)^2}\right\} \\
&= \mathcal{L}^{-1}\left\{\frac{s - \frac{3}{2} + \frac{3}{2}}{(s - \frac{3}{2})^2 + \left(\frac{\sqrt{3}}{2}\right)^2}\right\} \\
&= \mathcal{L}^{-1}\left\{\frac{s - \frac{3}{2}}{(s - \frac{3}{2})^2 + \left(\frac{\sqrt{3}}{2}\right)^2}\right\} + \mathcal{L}^{-1}\left\{\frac{\frac{3}{2}}{(s - \frac{3}{2})^2 + \left(\frac{\sqrt{3}}{2}\right)^2}\right\} \\
&= \mathcal{L}^{-1}\left\{\frac{s - \frac{3}{2}}{(s - \frac{3}{2})^2 + \left(\frac{\sqrt{3}}{2}\right)^2}\right\} + \sqrt{3} \mathcal{L}^{-1}\left\{\frac{\frac{\sqrt{3}}{2}}{(s - \frac{3}{2})^2 + \left(\frac{\sqrt{3}}{2}\right)^2}\right\} \\
&= e^{\frac{3}{2}t} \cos\left(\frac{\sqrt{3}}{2}t\right) + \sqrt{3} e^{\frac{3}{2}t} \sin\left(\frac{\sqrt{3}}{2}t\right)
\end{aligned}$$

SNB check:  $\frac{s}{s^2 - 3s + 3}$ , Is Laplace transform of  $e^{\frac{3}{2}t}(\cos \frac{1}{2}\sqrt{3}t + \sqrt{3} \sin \frac{1}{2}\sqrt{3}t)$

**3 (25 pts.)** Find the first six non-zero terms in the series solution near  $x = 0$  of the equation

$$y'' + 2xy' - 2y = 0$$

Be sure to give the recurrence relation. Indicate the two linearly independent solutions and give the first six nonzero terms of the solution.

Solution:

$$\begin{aligned}
y(x) &= \sum_{n=0}^{\infty} a_n x^n \\
y'(x) &= \sum_{n=1}^{\infty} a_n(n)x^{n-1} \\
y''(x) &= \sum_{n=2}^{\infty} a_n(n)(n-1)x^{n-2}
\end{aligned}$$

Substituting into the DE we have

$$\begin{aligned}
\sum_{n=2}^{\infty} a_n(n)(n-1)x^{n-2} + 2 \sum_{n=1}^{\infty} a_n(n)x^n - 2 \sum_{n=0}^{\infty} a_n x^n &= 0 \\
\sum_{n=2}^{\infty} a_n(n)(n-1)x^{n-2} + \sum_{n=1}^{\infty} a_n(2n-2)x^n - 2a_0 &= 0
\end{aligned}$$

Name: \_\_\_\_\_

Lecture Section \_\_\_\_

We now shift the first summation in the last equation above by letting  $k = n - 2$  or  $n = k + 2$ .

$$\sum_{k=0}^{\infty} a_{k+2}(k+2)(k+1)x^k + \sum_{n=1}^{\infty} a_n(2n-2)x^n - 2a_0 = 0$$

We write the first term of the first sum separately and replace  $k$  and  $n$  by  $m$  to get

$$\sum_{m=1}^{\infty} [a_{m+2}(m+2)(m+1) + a_m(2m-2)]x^m - 2a_0 + a_2(2)(1) = 0$$

Therefore

$$a_2 = a_0$$

and the recurrence relation is

$$a_{m+2}(m+2)(m+1) + a_m(2m-2) = 0 \quad m = 1, 2, \dots$$

or

$$a_{m+2} = \frac{2(1-m)}{(m+2)(m+1)} a_m \quad m = 1, 2, \dots$$

$$m = 1 \Rightarrow$$

$$a_3 = \frac{2(1-1)}{(3)(2)} a_1 = 0$$

$$m = 2 \Rightarrow$$

$$a_4 = \frac{2(1-2)}{(4)(3)} a_2 = \frac{2(-1)}{(4)(3)} a_0$$

$$m = 3 \Rightarrow$$

$$a_5 = \frac{2(1-3)}{(5)(4)} a_3 = 0$$

In fact all of the odd coefficients are zero except for  $a_1$ .

$$m = 4 \Rightarrow$$

$$a_6 = \frac{2(1-4)}{(6)(5)} a_4 = \frac{(2)(-3)}{(6)(5)} \frac{2(-1)}{(4)(3)} a_0 = \frac{(2)(2)(-3)(-1)}{(6)(5)(4)(3)} a_0$$

$$m = 6 \Rightarrow$$

$$a_8 = \frac{2(1-6)}{(8)(7)} a_6 = \frac{(2)(-5)}{(8)(7)} \frac{(2)(2)(3)(1)}{(6)(5)(4)(3)} a_0 = \frac{(2)(2)(2)(-5)(-3)(-1)}{(8)(7)(6)(5)(4)(3)} a_0$$

$$\begin{aligned} y(x) &= \sum_{n=0}^{\infty} a_n x^n = a_0 + a_1 x + a_2 x^2 + \dots \\ &= a_0 \left[ 1 + x^2 - \frac{2}{(4)(3)} x^4 + \frac{(2)(2)(3)(1)}{(6)(5)(4)(3)} x^6 - \frac{(2)(2)(2)(5)(3)(1)}{(8)(7)(6)(5)(4)(3)} x^8 + \dots \right] + a_1 x \end{aligned}$$

**4 (25 pts.)** Find the eigenvalues and eigenfunctions for

$$y'' + \lambda y = 0; \quad y(0) = 0, \quad y(\pi) = 0$$

Be sure to consider all values of  $\lambda$ .

Solution: The characteristic equation is:

$$\begin{aligned} p(r) &= r^2 + \lambda = 0 \\ \Rightarrow r &= \sqrt{-\lambda} \end{aligned}$$

There are three cases:

Case 1:  $-\lambda = 0$

$\Rightarrow r = 0$  is a repeated root.

$$\Rightarrow y = c_1 + c_2 x$$

$$0 = y(0) = c_1$$

$$0 = y(\pi) = c_2 \pi \Rightarrow c_2 = 0$$

$\Rightarrow y \equiv 0$  (trivial)

Case 2:  $-\lambda > 0$ , that is  $\lambda < 0$  Let  $\lambda = -k^2$  where  $k \neq 0$ .

$$r = \pm k$$

$$\Rightarrow y = c_1 e^{kx} + c_2 e^{-kx}$$

$$0 = y(0) = c_1 + c_2$$

$$\Rightarrow c_2 = -c_1$$

$$0 = y(\pi) = c_1 e^{k\pi} - c_1 e^{-k\pi}$$

$$0 = c_1 (e^{k\pi} - e^{-k\pi})$$

$\Rightarrow c_1 = 0 \Rightarrow c_2 = 0 \Rightarrow y \equiv 0$  (trivial)

Case 3:  $-\lambda < 0$ , that is  $\lambda > 0$  Let  $\lambda = k^2$  where  $k \neq 0$

$$\Rightarrow r = \pm ki$$

$$\Rightarrow y = c_1 \cos kx + c_2 \sin kx$$

$$0 = y(0) = c_1$$

$$0 = y(\pi) = c_2 \sin k\pi$$

$$\sin k\pi = 0 \Rightarrow k = n, \quad n = \dots, -3, -2, -1, 0, 1, 2, 3, \dots$$

$$\lambda_n = k^2 = n^2$$

$\Rightarrow$  The eigenvalues are:  $\lambda_n = n^2$  where  $n = 1, 2, 3, \dots$

and the eigenfunctions are:  $y_n = c_n \sin(nx)$  where  $n = 1, 2, 3, \dots$  and  $c_n$  are constants

Name: \_\_\_\_\_

Lecture Section \_\_\_\_

## Table of Laplace Transforms

$f(t)$	$F(s) = \mathcal{L}\{f\}(s)$		
$\frac{t^{n-1}}{(n-1)!}$	$\frac{1}{s^n}$	$n \geq 1$	$s > 0$
$e^{at}$	$\frac{1}{s-a}$		$s > a$
$\sin bt$	$\frac{b}{s^2 + b^2}$		$s > a$
$\cos bt$	$\frac{s}{s^2 + b^2}$		$s > a$
$e^{at}f(t)$	$\mathcal{L}\{f\}(s-a)$		
$t^n f(t)$	$(-1)^n \frac{d^n}{ds^n} (\mathcal{L}\{f\}(s))$		