

Name: \_\_\_\_\_

Lecturer \_\_\_\_\_

Lecture Section: \_\_\_\_\_

**Ma 221**

**Exam IA**

**15S**

**You may not use a calculator, cell phone, or computer while taking this exam. All work must be shown to obtain full credit. Credit will not be given for work not reasonably supported. When you finish, be sure to sign the pledge.**

Score on Problem #1 \_\_\_\_\_

#2 \_\_\_\_\_

#3 \_\_\_\_\_

#4 \_\_\_\_\_

Total Score \_\_\_\_\_

**I pledge my honor that I have abided by the Stevens Honor System.**

\_\_\_\_\_

Name: \_\_\_\_\_

Lecturer \_\_\_\_\_

Lecture Section: \_\_\_\_\_

**1 [ 20 pts. ]** Solve the initial value problem

$$\frac{dy}{dx} = \frac{2y}{x} + x^2 \cos x \quad y(\pi) = 2\pi^2.$$

Name: \_\_\_\_\_

Lecturer \_\_\_\_\_

Lecture Section: \_\_\_\_\_

**2 [ 20 pts. ]** Solve the initial value problem

$$2xy^3 dx - (1 - x^2) dy = 0 \quad y(0) = 1.$$

.

Name: \_\_\_\_\_

Lecturer \_\_\_\_\_

Lecture Section: \_\_\_\_\_

**3 [35 points]** Consider the differential equation

$$(3x^2y)dx + (3x^3 + 3)dy = 0$$

a. Show that the differential equation is not exact.

b. Find a value of  $n$ , such that multiplying the equation by  $y^n$  results in an exact differential equation.

c. The differential equation

$$(3x^2y^2 + 2x)dx + (2x^3y + 3y^2)dy = 0$$

is exact. Find a solution.

Name: \_\_\_\_\_ Lecturer \_\_\_\_\_

Lecture Section: \_\_\_\_\_

**4 [ 25 pts. ]**

$$\frac{dy}{dt} + 2y = y^2$$

Name: \_\_\_\_\_

Lecturer \_\_\_\_\_

Lecture Section: \_\_\_\_\_

## Table of Integrals

$\int \sec^2 t dt = \tan t + C$
$\int \frac{\sec^2 t}{\tan t} dt = \ln(\tan t) + C$
$\int \tan t dt = \ln(\sec t) + C$
$\int t e^{at} dt = \frac{1}{a^2} e^{at} (at - 1) + C$
$\int t^2 e^{at} dt = \frac{1}{a^3} e^{at} (a^2 t^2 - 2at + 2) + C$
$\int \cos^2 t dt = \frac{1}{2} t + \frac{1}{4} \sin 2t + C$
$\int \cos^3 t dt = \frac{1}{3} \cos^2 t \sin t + \frac{2}{3} \sin t + C$
$\int \sin^2 t dt = \frac{1}{2} t - \frac{1}{4} \pi - \frac{1}{4} \sin 2t + C$
$\int \sin^3 t dt = \frac{1}{12} \cos 3t - \frac{3}{4} \cos t + C$
$\int t \cos t dt = \cos t + t \sin t + C$
$\int t^2 \cos t dt = t^2 \sin t - 2 \sin t + 2t \cos t + C$
$\int t^3 \cos t dt = 3t^2 \cos t - 6 \cos t + t^3 \sin t - 6t \sin t + C$