

Name _____

Lecturer _____

Ma 221

Final Exam

5/7/09

Print Name: _____

Lecture Section: _____

I pledge my honor that I have abided by the Stevens Honor System.

This exam consists of 8 problems. You are to solve all of these problems. The point value of each problem is indicated. The total number of points is 200.

If you need more work space, continue the problem you are doing on the **other side of the page it is on**. Be sure that you do all problems.

You may not use a calculator, cell phone, or computer while taking this exam. All work must be shown to obtain full credit. Credit will not be given for work not reasonably supported. When you finish, be sure to sign the pledge.

There are tables giving Laplace transforms and integrals at the end of the exam.

Score on Problem #1 _____

#2 _____

#3 _____

#4 _____

#5 _____

#6 _____

#7 _____

#8 _____

Total Score _____

Name _____

Lecturer _____

1. Solve

(a) (8 pts)

$$x \cos x dx + (1 - 6y^5) dy = 0 \quad y(\pi) = 0$$

(b) (7 pts) Solve

$$y' = \frac{2 + ye^{xy}}{2y - xe^{xy}}$$

Name _____

Lecturer _____

1 (c) (10 pts) Find a general solution of

$$y'' - y' - 2y = 64e^{-t} + 4t^2$$

Name_____

Lecturer_____

2. (a) (12 pts) Find a general solution of

$$y'' - 2y' + y = 4 \sin x \quad x > 0$$

2(b) (13 pts.) Find a general solution of

$$x^2 y'' - xy' = x^3 e^x, \quad x > 0$$

Name _____

Lecturer _____

3. (a) (10 pts.) Let

$$g(t) = \begin{cases} e^t & \text{for } 0 \leq t \leq 2 \\ 3 & \text{for } 2 < t < \infty \end{cases}$$

Use the definition of the Laplace transform to find $\mathcal{L}\{g(t)\}$.

Name_____

Lecturer_____

(b) (15 pts.) Solve using Laplace Transforms:

$$y'' - 3y' + 4y = 0 \quad y(0) = 1, \quad y'(0) = 5$$

Name_____

Lecturer_____

4.) a.) (10 pts.) Use separation of variables, $u(x,t) = X(x)T(t)$, to find two ordinary differential equations which $X(x)$ and $T(t)$ must satisfy to be a solution of

$$5x^2t^7 \frac{\partial^3 u}{\partial x^2 \partial t} + (x+2)^3(t^2+10)^5 \frac{\partial^2 u}{\partial t^2} = 0$$

Note: Do **not** solve these ordinary differential equations.

b.) (15 pts.) Find the eigenvalues and eigenfunctions for Find the eigenvalues and eigenfunctions for

$$y'' - 4\lambda y' + 4\lambda^2 y = 0 \quad y'(1) = 0, \quad y(2) + 2y'(2) = 0$$

Name _____

Lecturer _____

5. (a) (15 pts.) Find the first four nonzero terms of the Fourier *sine* series for the function

$$f(x) = \begin{cases} x & ; \quad 0 < x < \pi/2 \\ 0 & ; \quad \pi/2 < x < \pi \end{cases}$$

Be sure to give the Fourier series with these terms in it.

(b) (10 pts.) Sketch the graph of the function represented by the Fourier sine series in 5 (a) on $-\pi < x < 3\pi$.

Name _____

Lecturer _____

6 (25 pts.)

PDE $u_t = 3u_{xx}$

BCs $u(0, t) = 0 \quad u_x(2, t) = 0$

IC $u(x, 0) = 2 \sin\left(\frac{3\pi}{4}x\right) - 7 \sin\left(\frac{5\pi}{4}x\right)$

You must derive the solution. Your solution should not have any arbitrary constants in it. Show **all** steps.

Name _____

Lecturer _____

7. (a) (10 pts.) Solve

$$y'' - 4y' + 13y = 0; \quad y(0) = 0, \quad y(\pi) = 0$$

(b) (15 pts.) Find the first 6 nonzero terms of the power series solution about $x = 0$ for the DE:

$$y'' + 2x^2y' + 2xy = 0$$

Be sure to give the recurrence relation.

Name_____

Lecturer_____

8 (a) (12 pts.) Solve

$$x \frac{dy}{dx} = 3y + x^2 y^{-3}$$

(b) (13 pts.) Find

$$\mathcal{L}^{-1} \left\{ \frac{s^2 + s + 6}{s(s-1)(s-3)} \right\}$$

Table of Laplace Transforms

$f(t)$	$F(s) = \mathcal{L}\{f\}(s)$		
$\frac{t^{n-1}}{(n-1)!}$	$\frac{1}{s^n}$	$n \geq 1$	$s > 0$
e^{at}	$\frac{1}{s-a}$		$s > a$
$\sin bt$	$\frac{b}{s^2 + b^2}$		$s > 0$
$\cos bt$	$\frac{s}{s^2 + b^2}$		$s > 0$
$e^{at}f(t)$	$\mathcal{L}\{f\}(s-a)$		
$t^n f(t)$	$(-1)^n \frac{d^n}{ds^n} (\mathcal{L}\{f\}(s))$		

Table of Integrals

$\int \sin^2 x dx = -\frac{1}{2} \cos x \sin x + \frac{1}{2} x + C$
$\int \cos^2 x dx = \frac{1}{2} \cos x \sin x + \frac{1}{2} x + C$
$\int x \cos bx dx = \frac{1}{b^2} (\cos bx + bx \sin bx) + C$
$\int x \sin bx = \frac{1}{b^2} (\sin bx - bx \cos bx) + C$
$\int \left(\frac{e^{-t}}{1+e^t} \right) dt = -e^{-t} - \ln(e^t) + \ln(1+e^t) + C$
$\int x e^{ax} dx = \frac{1}{a^2} (axe^{ax} - e^{ax}) + C$