

Later we will say this again in different words: If every column of  $A$  is perpendicular to  $\mathbf{y} = (2, 1, -1)$ , then any combination  $\mathbf{b}$  of those columns must also be perpendicular to  $\mathbf{y}$ . Otherwise  $\mathbf{b}$  is not in the column space and  $A\mathbf{x} = \mathbf{b}$  is not solvable.

And again: If  $\mathbf{y}$  is in the nullspace of  $A^T$  then  $\mathbf{y}$  must be perpendicular to every  $\mathbf{b}$  in the column space. Just looking ahead ...

### Problem Set 3.4

- 1 (Recommended) Execute the six steps of Worked Example 3.4 A to describe the column space and nullspace of  $A$  and the complete solution to  $A\mathbf{x} = \mathbf{b}$ :

$$A = \begin{bmatrix} 2 & 4 & 6 & 4 \\ 2 & 5 & 7 & 6 \\ 2 & 3 & 5 & 2 \end{bmatrix} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \\ 5 \end{bmatrix}$$

- 2 Carry out the same six steps for this matrix  $A$  with rank one. You will find *two* conditions on  $b_1, b_2, b_3$  for  $A\mathbf{x} = \mathbf{b}$  to be solvable. Together these two conditions put  $\mathbf{b}$  into the \_\_\_\_\_ space (two planes give a line):

$$A = \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix} [2 \ 1 \ 3] = \begin{bmatrix} 2 & 1 & 3 \\ 6 & 3 & 9 \\ 4 & 2 & 6 \end{bmatrix} \quad \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 10 \\ 30 \\ 20 \end{bmatrix}$$

Questions 3–15 are about the solution of  $A\mathbf{x} = \mathbf{b}$ . Follow the steps in the text to  $\mathbf{x}_p$  and  $\mathbf{x}_n$ . Use the augmented matrix with last column  $\mathbf{b}$ .

- 3 Write the complete solution as  $\mathbf{x}_p$  plus any multiple of  $s$  in the nullspace:

$$\begin{aligned} x + 3y + 3z &= 1 \\ 2x + 6y + 9z &= 5 \\ -x - 3y + 3z &= 5. \end{aligned}$$

- 4 Find the complete solution (also called the *general solution*) to

$$\begin{bmatrix} 1 & 3 & 1 & 2 \\ 2 & 6 & 4 & 8 \\ 0 & 0 & 2 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix}.$$

- 5 Under what condition on  $b_1, b_2, b_3$  is this system solvable? Include  $\mathbf{b}$  as a fourth column in elimination. Find all solutions when that condition holds:

$$\begin{aligned} x + 2y - 2z &= b_1 \\ 2x + 5y - 4z &= b_2 \\ 4x + 9y - 8z &= b_3. \end{aligned}$$

- 6 What conditions on  $b_1, b_2, b_3, b_4$  make each system solvable? Find  $x$  in that case:

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \\ 2 & 5 \\ 3 & 9 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix} \qquad \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 2 & 5 & 7 \\ 3 & 9 & 12 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}$$

- 7 Show by elimination that  $(b_1, b_2, b_3)$  is in the column space if  $b_3 - 2b_2 + 4b_1 = 0$ .

$$A = \begin{bmatrix} 1 & 3 & 1 \\ 3 & 8 & 2 \\ 2 & 4 & 0 \end{bmatrix}$$

What combination of the rows of  $A$  gives the zero row?

- 8 Which vectors  $(b_1, b_2, b_3)$  are in the column space of  $A$ ? Which combinations of the rows of  $A$  give zero?

(a)  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 6 & 3 \\ 0 & 2 & 5 \end{bmatrix}$

(b)  $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 2 & 4 & 8 \end{bmatrix}$

- 9 (a) The Worked Example 3.4  $A$  reached  $[U \ c]$  from  $[A \ b]$ . Put the multipliers into  $L$  and verify that  $LU$  equals  $A$  and  $Lc$  equals  $b$ .  
 (b) Combine the pivot columns of  $A$  with the numbers  $-9$  and  $3$  in the particular solution  $x_p$ . What is that linear combination and why?
- 10 Construct a 2 by 3 system  $Ax = b$  with particular solution  $x_p = (2, 4, 0)$  and homogeneous solution  $x_n =$  any multiple of  $(1, 1, 1)$ .
- 11 Why can't a 1 by 3 system have  $x_p = (2, 4, 0)$  and  $x_n =$  any multiple of  $(1, 1, 1)$ ?
- 12 (a) If  $Ax = b$  has two solutions  $x_1$  and  $x_2$ , find two solutions to  $Ax = 0$ .  
 (b) Then find another solution to  $Ax = 0$  and another solution to  $Ax = b$ .
- 13 Explain why these are all false:  
 (a) The complete solution is any linear combination of  $x_p$  and  $x_n$ .  
 (b) A system  $Ax = b$  has at most one particular solution.  
 (c) The solution  $x_p$  with all free variables zero is the shortest solution (minimum length  $\|x\|$ ). Find a 2 by 2 counterexample.  
 (d) If  $A$  is invertible there is no solution  $x_n$  in the nullspace.
- 14 Suppose column 5 of  $U$  has no pivot. Then  $x_5$  is a \_\_\_\_\_ variable. The zero vector (is) (is not) the only solution to  $Ax = 0$ . If  $Ax = b$  has a solution, then it has \_\_\_\_\_ solutions.