## Assignment 11.

This homework is due *Tuesday* Nov 29.

There are total 48 points in this assignment. 43 points is considered 100%. If you go over 43 points, you will get over 100% for this homework (up to 115%) and it will count towards your course grade.

Collaboration is welcome. If you do collaborate, make sure to write/type your own paper and give credit to your collaborators in your pledge. Your solutions should contain full proofs. Bare answers will not earn you much.

This assignment covers sections 6.2, 6.4 in Bartle–Sherbert.

- (1) [3pt] (Exercise 6.2.3a) Find the points of relative extrema of the function  $f(x) = |x^2 1|$  for  $-4 \le x \le 4$ .
- (2) [3pt] (Exercise 6.2.6) Prove that  $|\sin x \sin y| \le |x y|$  for all  $x, y \in \mathbb{R}$ . (Hint: Apply the Mean Value theorem to sin on the interval [x, y].)
- (3) (a) [3pt] (Exercise 6.2.8) Let  $f : [a,b] \to \mathbb{R}$  be continuous on [a,b] and differentiable on (a,b). Show that if  $\lim_{x\to a} f'(x) = A$ , then f'(a) exists and is equal to A. (Hint: Use the limit definition of f'(a) and apply the Mean Value Theorem to f on the interval [a, x].)
  - (b) [3pt] Is function

$$f(x) = \begin{cases} \frac{1}{x} - \frac{1}{e^x - 1}, & x \neq 0; \\ 1/2, & x = 0, \end{cases}$$

differentiable at 0?

- (4) [3pt] (Exercise 6.2.17) Let f, g be differentiable on  $\mathbb{R}$  and suppose that f(0) = g(0), and  $f'(x) \leq g'(x)$  for all  $x \geq 0$ . Show that  $f(x) \leq g(x)$  for all  $x \geq 0$ . (Hint: Apply the Mean Value Theorem to f g on [0, x].)
- (5) For a given function f and a point  $x_0$ , find Taylor's polynomials  $P_2(x)$ ,  $P_3(x)$ ,  $P_4(x)$ ,  $P_5(x)$ ,  $P_{2011}(x)$  of f(x) at  $x_0 = \pi/2$ .
  - (a) [2pt]  $f(x) = \sin x$  at  $x_0 = \pi/2$ . Compare to  $\cos at 0$ .
  - (b) [2pt]  $f(x) = \cos x$  at  $x_0 = -\pi/2$ . Compare to sin at 0.
  - (c) [2pt]  $f(x) = x^3$  at  $x_0 = 2$ . Compare  $P_3(x), P_4(x), P_{2011}(x)$  to f(x).
  - (d) [2pt]  $f(x) = \frac{1}{1-x}$  at  $x_0 = 0$ .
  - (e) [2pt]  $f(x) = \frac{1}{x}$  at  $x_0 = 1$ . Compare to the previous item.
- (6) [4pt] (Part of exercise 6.4.7) If x > 0, show that

$$\sqrt[3]{1+x} - \left(1 + \frac{1}{3}x - \frac{1}{9}x^2\right) \le \frac{5}{81}x^3.$$

(*Hint:* Use Taylor's Theorem with n = 2.)

- see next page -

(7) (a) [3pt] Suppose  $A \in \mathbb{R}$ . Show that  $\lim_{n \to \infty} \frac{A^n}{n!} = 0$ . *Hint:* take tail of this sequence that starts with m > 2|A| and represent

$$\frac{A^n}{n!} = \frac{A^m}{m!} \cdot \frac{A^{n-m}}{(m+1)\cdots n}.$$

- (b) [4pt] (Exercise 6.4.8) If  $f(x) = e^x$ , show that the remainder term in Taylor's Theorem converges to zero as  $n \to \infty$ , for each fixed  $x_0$  and x.
- (c) [4pt] (Exercise 6.4.9) If  $g(x) = \sin x$ , show that the remainder term in Taylor's Theorem converges to zero as  $n \to \infty$ , for each fixed  $x_0$  and x.
- (8) (Part of exercise 6.4.14) Determine whether or not x = 0 is a point of relative extremum of the following functions:
  (a) [2pt] f(x) = x<sup>3</sup> + 2,
  - (a) [2pt] f(x) = x + 2,
  - (b) [3pt]  $f(x) = \sin x x$ ,
  - (c) [3pt]  $f(x) = \cos x 1 + \frac{1}{2}x^2$ .

 $\mathbf{2}$